Contents

Volume 121 1995

Felsenfeld, A. L. and Kennison, J. A.
Positional signaling by *hedgehog* in *Drosophila* imaginal disc development 1-10

Guz, Y., Montminy, M. R., Stein, R., Leonard, J., Gamer, L. W., Wright, C. V. E. and Teitelman, G.
Expression of murine STF-1, a putative insulin gene transcription factor, in β cells of pancreas, duodenal epithelium and pancreatic exocrine and endocrine progenitors during ontogeny 11-18

Carpenter, R. and Coen, E. S.
Transposon induced chimeras show that *floricaula*, a meristem identity gene, acts non-autonomously between cell layers 19-26

Hantke, S. S., Carpenter, R. and Coen, E. S.
Expression of *floricaula* in single cell layers of periclinal chimeras activates downstream homeotic genes in all layers of floral meristems 27-35

Mitsiadis, T. A., Salmivirta, M., Muramatsu, T., Muramatsu, H., Rauvala, H., Lehtonen, E., Jalkanen, M. and Thesleff, I.
Expression of the heparin-binding cytokines, midkine (MK) and HB-GAM (pleiotrophin) is associated with epithelial-mesenchymal interactions during fetal development and organogenesis 37-51

Scheres, B., Di Laurenzio, L., Willemsen, V., Hauser, M.-T., Janmaat, K., Weisbeek, P. and Benfey, P. N.
Mutations affecting the radial organisation of the *Arabidopsis* root display specific defects throughout the embryonic axis 53-62

Ruden, D. M. and Jäckle, H.
Mitotic delay dependent survival identifies components of cell cycle control in the *Drosophila* blastoderm 63-73

Boyan, G., Therianos, S., Williams, J. L. D. and Reichert, H.
Axogenesis in the embryonic brain of the grasshopper *Schistocerca gregaria*: an identified cell analysis of early brain development 75-86

Quinlan, G. A., Williams, E. A., Tan, S.-S. and Tam, P. P. L.
Neuroectodermal fate of epiblast cells in the distal region of the mouse egg cylinder: implication for body plan organization during early embryogenesis 87-98

Pultz, M. A. and Baker, B. S.
The dual role of *hermaphrodite* in the *Drosophila* sex determination regulatory hierarchy 99-111

Christians, E., Campion, E., Thompson, E. M. and Renard, J.-P.
Expression of the *HSP 70.1* gene, a landmark of early zygotic activity in the mouse embryo, is restricted to the first burst of transcription 113-122

Ganfornina, M. D., Sánchez, D. and Bastiani, M. J.
Lazarillo, a new GPI-linked surface lipocalin, is restricted to a subset of neurons in the grasshopper embryo 123-134

Sánchez, D., Ganfornina, M. D. and Bastiani, M. J.
Developmental expression of the lipocalin Lazarillo and its role in axonal pathfinding in the grasshopper embryo 135-147

Sharp, R., Babyatsky, M. W., Takagi, H., Tågerud, S., Wang, T. C., Bockman, D. E., Brand, S. J. and Merlino, G.
Transforming growth factor α disrupts the normal program of cellular differentiation in the gastric mucosa of transgenic mice 149-161

Pevny, L., Lin, C.-S., D'Agati, V., Simon, M. C., Orkin, S. H. and Costantini, F.
Development of hematopoietic cells lacking transcription factor GATA-1 163-172

Godt, D. and Laski, F. A.
Mechanisms of cell rearrangement and cell recruitment in *Drosophila* ovary morphogenesis and the requirement of *bric à brac* 173-187

Yamagata, M. and Sanes, J. R.
Lamina-specific cues guide outgrowth and arborization of retinal axons in the optic tectum 189-200

Forristall, C., Pondel, M., Chen, L. and King, M. L.
Patterns of localization and cytoskeletal association of two vegetally localized RNAs, *Vg1* and *Xcat-2* 201-208

Graba, Y., Gieseler, K., Aragnol, D., Laurenti, P., Mariol, M.-C., Berenger, H., Sagnier, T. and Pradel, J.
DWnt-4, a novel *Drosophila Wnt* gene acts downstream of homeotic complex genes in the visceral mesoderm 209-218

Martín-Bermudo, M. D., Carmena, A. and Jiménez, F.
Neurogenic genes control gene expression at the transcriptional level in early neurogenesis and in mesectoderm specification 219-224

Begemann, G., Michon, A.-M., Voorn, L. v.d., Wepf, R. and Mlodzik, M.
The *Drosophila* orphan nuclear receptor Seven-up requires the Ras pathway for its function in photoreceptor determination 225-235

Mohler, J., Mahaffey, J. W., Deutsch, E. and Vani, K.
Control of *Drosophila* head segment identity by the bZIP homeotic gene *cnc* 237-247

Tucker, A. S. and Slack, J. M. W.
The *Xenopus laevis* tail-forming region 249-262

Newman, A. P., White, J. G. and Sternberg, P. W.
The *Caenorhabditis elegans lin-12* gene mediates induction of ventral uterine specialization by the anchor cell 263-271

Faust, C., Schumacher, A., Holdener, B. and Magnuson, T.
The *eed* mutation disrupts anterior mesoderm production in mice 273-285

Kloc, M. and Etkin, L. D.
Two distinct pathways for the localization of RNAs at the vegetal cortex in *Xenopus* oocytes 287-297

Chia, C. M., Winston, R. M. L. and Handyside, A. H.
EGF, TGF-α and EGFR expression in human preimplantation embryos 299-307

Ekström, T. J., Cui, H., Li, X. and Ohlsson, R.
Promoter-specific *IGF2* imprinting status and its plasticity during human liver development 309-316

Halter, D. A., Urban, J., Rickert, C., Ner, S. S., Ito, K., Travers, A. A. and Technau, G. M.
The homeobox gene *repo* is required for the differentiation and maintenance of glia function in the embryonic nervous system of *Drosophila melanogaster* 317-332

Burke, A. C., Nelson, C. E., Morgan, B. A. and Tabin, C.
Hox genes and the evolution of vertebrate axial morphology 333-346

Akimenko, M.-A., Johnson, S. L., Westerfield, M. and Ekker, M.
Differential induction of four *msx* homeobox genes during fin development and regeneration in zebrafish 347-357

Becker, T., Berliner, A. J., Nitabach, M. N., Gan, W.-B. and Macagno, E. R.
Target-induced neurogenesis in the leech CNS involves efferent projections to the target 359-369

Paul, D. L., Yu, K., Bruzzone, R., Gimlich, R. L. and Goodenough, D. A.
Expression of a dominant negative inhibitor of intercellular communication in the early *Xenopus* embryo causes delamination and extrusion of cells 371-381

Toyama, R., O'Connell, M. L., Wright, C. V. E., Kuehn, M. R. and Dawid, I. B.
Nodal induces ectopic *goosecoid* and *lim1* expression and axis duplication in zebrafish 383-391

Tepass, U. and Hartenstein, V.
Neurogenic and proneural genes control cell fate specification in the *Drosophila* endoderm 393-405

Downs, K. M. and Gardner, R. L.
An investigation into early placental ontogeny: allantoic attachment to the chorion is selective and developmentally regulated 407-416

Storey, K. G., Selleck, M. A. J. and Stern, C. D.
Neural induction and regionalisation by different subpopulations of cells in Hensen's node 417-428

Giangrande, A.
Proneural genes influence gliogenesis in *Drosophila* 429-438

Crossley, P. H. and Martin, G. R.
The mouse *Fgf8* gene encodes a family of polypeptides and is expressed in regions that direct outgrowth and patterning in the developing embryo 439-451

Tsai, C. and Gergen, P.
Pair-rule expression of the *Drosophila fushi tarazu* gene: a nuclear receptor response element mediates the opposing regulatory effects of *runt* and *hairy* 453-462

Gorman, M., Franke, A. and Baker, B. S.
Molecular characterization of the *male-specific lethal-3* gene and investigations of the regulation of dosage compensation in *Drosophila* 463-475

Wilder, E. L. and Perrimon, N.
Dual functions of *wingless* in the *Drosophila* leg imaginal disc 477-488

Kwee, L., Baldwin, H. S., Shen, H. M., Stewart, C. L., Buck, C., Buck, C. A. and Labow, M. A.
Defective development of the embryonic and extraembryonic circulatory systems in vascular cell adhesion molecule (VCAM-1) deficient mice 489-503

Robinson, M. L., Overbeek, P. A., Verran, D. J., Grizzle, W. E., Stockard, C. R., Friesel, R., Maciag, T. and Thompson, J. A.
Extracellular FGF-1 acts as a lens differentiation factor in transgenic mice 505-514

Nascone, N. and Mercola, M.
An inductive role for the endoderm in *Xenopus* cardiogenesis 515-523

Selleck, M. A. J. and Bronner-Fraser, M.
Origins of the avian neural crest: the role of neural plate-epidermal interactions 525-538

Lampron, C., Rochette-Egly, C., Gorry, P., Dollé, P., Mark, M., Lufkin, T., LeMeur, M. and Chambon, P.
Mice deficient in cellular retinoic acid binding protein II (CRABPII) or in both CRABPI and CRABPII are essentially normal 539-548

Yang, J. T., Rayburn, H. and Hynes, R. O.
Cell adhesion events mediated by α_4 integrins are essential in placental and cardiac development 549-560

O'Neill, M. J. and Artzt, K.
Identification of a germ-cell-specific transcriptional repressor in the promoter of *Tctex-1* 561-568

Sonnenfeld, M. J. and Jacobs, J. R.
Apoptosis of the midline glia during *Drosophila* embryogenesis: a correlation with axon contact 569-578

Nagoshi, R. N., Patton, J. S., Bae, E. and Geyer, P. K.
The somatic sex determines the requirement for *ovarian tumor* gene activity in the proliferation of the *Drosophila* germline 579-587

Ng, M., Diaz-Benjumea, F. J. and Cohen, S. M.
nubbin encodes a POU-domain protein required for proximal-distal patterning in the *Drosophila* wing 589-599

Ferreira, J. and Carmo-Fonseca, M.
The biogenesis of the coiled body during early mouse development 601-612

Submission of sequence data to the EMBL nucleotide sequence database 613-618

ESSAY IN DEVELOPMENT

Campbell, G. and Tomlinson, A.
Initiation of the proximodistal axis in insect legs 619-628

CONTRIBUTED PAPERS

Nambara, E., Keith, K., McCourt, P. and Naito, S.
A regulatory role for the *ABI3* gene in the establishment of embryo maturation in *Arabidopsis thaliana* 629-636

Goldhamer, D. J., Brunk, B. P., Faerman, A., King, A., Shani, M. and Emerson, C. P., Jr
Embryonic activation of the *myoD* gene is regulated by a highly conserved distal control element 637-649

Münsterberg, A. E. and Lassar, A. B.
Combinatorial signals from the neural tube, floor plate and notochord induce myogenic bHLH gene expression in the somite 651-660

Hayashi, K. and Ozawa, E.
Myogenic cell migration from somites is induced by tissue contact with medial region of the presumptive limb mesoderm in chick embryos 661-669

Fawcett, D., Pasceri, P., Fraser, R., Colbert, M., Rossant, J. and Giguère, V.
Postaxial polydactyly in forelimbs of *CRABP-II* mutant mice 671-679

Chen, W.-H., Morriss-Kay, G. M. and Copp, A. J.
Genesis and prevention of spinal neural tube defects in the *curly tail* mutant mouse: involvement of retinoic acid and its nuclear receptors RAR-β and RAR-γ 681-691

Oliver, G., Wehr, R., Jenkins, N. A., Copeland, N. G., Cheyette, B. N. R., Hartenstein, V., Zipursky, S. L. and Gruss, P.
Homeobox genes and connective tissue patterning 693-705

Pannese, M., Polo, C., Andreazzoli, M., Vignali, R., Kablar, B., Barsacchi, G. and Boncinelli, E.
The *Xenopus* homologue of *Otx2* is a maternal homeobox gene that demarcates and specifies anterior body regions 707-720

Witta, S. E., Agarwal, V. R. and Sato, S. M.
XIPOU 2, a noggin-inducible gene, has direct neuralizing activity 721-730

Wehrle-Haller, B. and Weston, J. A.
Soluble and cell-bound forms of steel factor activity play distinct roles in melanocyte precursor dispersal and survival on the lateral neural crest migration pathway 731-742

Collins, J. E., Lorimer, J. E., Garrod, D. R., Pidsley, S. C., Buxton, R. S. and Fleming, T. P.
Regulation of desmocollin transcription in mouse preimplantation embryos 743-753

Pierce, S. B. and Kimelman, D.
Regulation of Spemann organizer formation by the intracellular kinase Xgsk-3 755-765

Mayor, R., Morgan, R. and Sargent, M. G.
Induction of the prospective neural crest of *Xenopus* 767-777

Kato, Y. and Tsunoda, Y.
Germ cell nuclei of male fetal mice can support development of chimeras to midgestation following serial transplantation 779-783

Sturtevant, M. A. and Bier, E.
Analysis of the genetic hierarchy guiding wing vein development in *Drosophila* 785-801

Fishell, G.
Striatal precursors adopt cortical identities in response to local cues 803-812

Streit, A., Stern, C. D., Théry, C., Ireland, G. W., Aparicio, S., Sharpe, M. J. and Gherardi, E.
A role for HGF/SF in neural induction and its expression in Hensen's node during gastrulation 813-824

Lee, Y. M., Osumi-Yamashita, N., Ninomiya, Y., Moon, C. K., Eriksson, U. and Eto, K.
Retinoic acid stage-dependently alters the migration pattern and identity of hindbrain neural crest cells 825-837

Monaghan, A. P., Grau, E., Bock, D. and Schütz, G.
The mouse homolog of the orphan nuclear receptor *tailless* is expressed in the developing forebrain 839-853

Gu, Y., Hukriede, N. A. and Fleming, R. J.
Serrate expression can functionally replace *Delta* activity during neuroblast segregation in the *Drosophila* embryo 855-865

Ryan, G., Steele-Perkins, V., Morris, J. F., Rauscher III, F. J. and Dressler, G. R.
Repression of *Pax-2* by *WT1* during normal kidney development 867-875

Wilson, V., Manson, L., Skarnes , W. C. and Beddington, R. S. P.
The *T* gene is necessary for normal mesodermal morphogenetic cell movements during gastrulation 877-886

Groves, A. K., George, K. M., Tissier-Seta, J.-P., Engel, J. D., Brunet, J.-F. and Anderson, D. J.
Differential regulation of transcription factor gene expression and phenotypic markers in developing sympathetic neurons 887-901

Harden, N., Loh, H. Y., Chia, W. and Lim, L.
A dominant inhibitory version of the small GTP-binding protein Rac disrupts cytoskeletal structures and inhibits developmental cell shape changes in *Drosophila* 903-914

Erickson, C. A. and Goins, T. L.
Avian neural crest cells can migrate in the dorsolateral path only if they are specified as melanocytes 915-924

Hampl, A. and Eppig, J. J.
Analysis of the mechanism(s) of metaphase I arrest in maturing mouse oocytes 925-933

Birgbauer, E., Sechrist, J., Bronner-Fraser, M. and Fraser, S.
Rhombomeric origin and rostrocaudal reassortment of neural crest cells revealed by intravital microscopy 935-945

Lefebvre, O., Régnier, C., Chenard, M.-P., Wendling, C., Chambon, P., Basset , P. and Rio, M.-C.
Developmental expression of mouse stromelysin-3 mRNA 947-955

Frasch, M., Chen, X. and Lufkin, T.
Evolutionary-conserved enhancers direct region-specific expression of the murine *Hoxa-1* and *Hoxa-2* loci in both mice and *Drosophila* 957-974

Liu, Z. and Meyerowitz, E. M.
LEUNIG regulates *AGAMOUS* expression in *Arabidopsis* flowers 975-991

Blitz, I. L. and Cho, K. W. Y.
Anterior neurectoderm is progressively induced during gastrulation: the role of the *Xenopus* homeobox gene *orthodenticle* 993-1004

Harvey, M. B., Leco, K. J., Arcellana-Panlilio, M. Y., Zhang, X., Edwards, D. R. and Schultz, G. A.
Proteinase expression in early mouse embryos is regulated by leukaemia inhibitory factor and epidermal growth factor 1005-1014

Nogawa, H. and Ito, T.
Branching morphogenesis of embryonic mouse lung epithelium in mesenchyme-free culture 1015-1022

Schulz, C. and Tautz, D.
Zygotic *caudal* regulation by *hunchback* and its role in abdominal segment formation of the *Drosophila* embryo 1023-1028

Tan, S.-S., Faulkner-Jones, B., Breen, S. J., Walsh, M., Bertram, J. F. and Reese, B. E.
Cell dispersion patterns in different cortical regions studied with an X-inactivated transgenic marker 1029-1039

Cary, R. B. and Klymkowsky, M. W.
Disruption of intermediate filament organization leads to structural defects at the intersomite junction in *Xenopus* myotomal muscle 1041-1052

Xu, T., Wang, W., Zhang, S., Stewart, R. A. and Yu, W.
Identifying tumor suppressors in genetic mosaics: the *Drosophila lats* gene encodes a putative protein kinase 1053-1063

Reginelli, A. D., Wang, Y.-Q., Sassoon, D. and Muneoka, K.
Digit tip regeneration correlates with regions of *Msx1* (*Hox 7*) expression in fetal and newborn mice 1065-1076

Knittel, T., Kessel, M., Kim, M. H. and Gruss, P.
A conserved enhancer of the human and murine *Hoxa-7* gene specifies the anterior boundary of expression during embryonal development 1077-1088

Schlaeger, T. M., Qin, Y., Fujiwara, Y., Magram, J. and Sato, T. N.
Vascular endothelial cell lineage-specific promoter in transgenic mice 1089-1098

Cserjesi, P., Brown, D., Ligon, K. L., Lyons, G. E., Copeland, N. G., Gilbert, D. J., Jenkins, N. A. and Olson, E. N.
Scleraxis: a basic helix-loop-helix protein that prefigures skeletal formation during mouse embryogenesis 1099-1110

Wang, D. G.-W., Kirchhamer, C. V., Britten, R. J. and Davidson, E. H.
SpZ12-1, a negative regulator required for spatial control of the territory-specific *CyIIIa* gene in the sea urchin embryo 1111-1122

Kono, T., Carroll, J., Swann, K. and Whittingham, D. G.
Nuclei from fertilized mouse embryos have calcium-releasing activity 1123-1128

Visconti, P. E., Bailey, J. L., Moore, G. D., Pan, D., Olds-Clarke, P. and Kopf, G. S.
Capacitation of mouse spermatozoa. I. Correlation between the capacitation state and protein tyrosine phosphorylation 1129-1137

Visconti, P. E., Moore, G. D., Bailey, J. L., Leclerc, P., Connors, S. A., Pan, D., Olds-Clarke, P. and Kopf, G. S.
Capacitation of mouse spermatozoa. II. Protein tyrosine phosphorylation and capacitation are regulated by a cAMP-dependent pathway 1139-1150

Ferri, R. T. and Levitt, P.
Regulation of regional differences in the differentiation of cerebral cortical neurons by EGF family-matrix interactions 1151-1160

Zhukareva, V. and Levitt, P.
The limbic system-associated membrane protein (LAMP) selectively mediates interactions with specific central neuron populations 1161-1172

Datta, S.
Control of proliferation activation in quiescent neuroblasts of the *Drosophila* central nervous system 1173-1182

Norman, D. J., Feng, L., Cheng, S. S., Gubbay, J., Chan, E. and Heintz, N.
The lurcher gene induces apoptotic death in cerebellar Purkinje cells 1183-1193

Huet, F., Ruiz, C. and Richards, G.
Sequential gene activation by ecdysone in *Drosophila melanogaster*: the hierarchical equivalence of early and early late genes 1195-1204

Galliot, B., Welschof, M., Schuckert, O., Hoffmeister, S. and Schaller, H. C.
The cAMP Response Element Binding protein is involved in hydra regeneration 1205-1216

Pogge v. Strandmann, E. and Ryffel, G. U.
Developmental expression of the maternal protein XDCoH, the dimerization cofactor of the homeoprotein LFB1 (HNF1) 1217-1226

Goldstein, B.
An analysis of the response to gut induction in the *C. elegans* embryo 1227-1236

Hauser, M.-T., Morikami, A. and Benfey, P. N.
Conditional root expansion mutants of *Arabidopsis* 1237-1252

Svendsen, P. C. and McGhee, J. D.
The *C. elegans* neuronally expressed homeobox gene *ceh-10* is closely related to genes expressed in the vertebrate eye 1253-1262

Maves, L. and Schubiger, G.
wingless induces transdetermination in developing *Drosophila* imaginal discs 1263-1272

Technau, U. and Holstein, T. W.
Head formation in *Hydra* is different at apical and basal levels 1273-1282

Ware, C. B., Horowitz, M. C., Renshaw, B. R., Hunt, J. S., Liggitt, D., Koblar, S. A., Gliniak, B. C., McKenna, H. J., Papayannopoulou, T., Thoma, B., Cheng, L., Donovan, P. J., Peschon, J. J., Bartlett, P. F., Willis, C. R., Wright, B. D., Carpenter, M. K., Davison, B. L. and Gearing, D. P.
Targeted disruption of the low-affinity leukemia inhibitory factor receptor gene causes placental, skeletal, neural and metabolic defects and results in perinatal death 1283-1299

Tuttle, R. and Matthew, W. D.
Neurotrophins affect the pattern of DRG neurite growth in a bioassay that presents a choice of CNS and PNS substrates 1301-1309

Brower, D. L., Bunch, T. A., Mukai, L. Adamson, T. E., Wehrli, M., Lam, S., Friedlander, E., Roote, C. E. and Zusman, S.
Nonequivalent requirements for PS1 and PS2 integrin at cell attachments in *Drosophila*: genetic analysis of the α_{PS1} integrin subunit 1311-1320

Nakagawa, S. and Takeichi, M.
Neural crest cell-cell adhesion controlled by sequential and subpopulation-specific expression of novel cadherins 1321-1332

Bhatnagar, P., Papaioannou, V. E. and Biggers, J. D.
CSF-1 and mouse preimplantation development in vitro 1333-1339

Oakley, R. A., Garner, A. S., Large, T. H. and Frank, E.
Muscle sensory neurons require neurotrophin-3 from peripheral tissues during the period of normal cell death 1341-1350

O'Reilly, M.-A. J., Smith, J. C. and Cunliffe, V.
Patterning of the mesoderm in *Xenopus*: dose-dependent and synergistic effects of *Brachyury* and *Pintallavis* 1351-1359

Kramer, S., West, S. R. and Hiromi, Y.
Cell fate control in the *Drosophila* retina by the orphan receptor seven-up: its role in the decisions mediated by the ras signaling pathway 1361-1372

Hsieh-Li, H. M., Witte, D. P., Weinstein, M., Branford, W., Li, H., Small, K. and Potter, S. S.
Hoxa 11 structure, extensive antisense transcription, and function in male and female fertility 1373-1385

Baker, R. and Schubiger, G.
Ectoderm induces muscle-specific gene expression in *Drosophila* embryos 1387-1398

Mahmood, R., Kiefer, P., Guthrie, S., Dickson, C. and Mason, I.
Multiple roles for FGF-3 during cranial neural development in the chicken 1399-1410

Fletcher, J. C. and Thummel, C. S.
The *Drosophila E74* gene is required for the proper stage- and tissue-specific transcription of ecdysone-regulated genes at the onset of metamorphosis 1411-1421

Gourdie, R. G., Mima, T., Thompson, R. P. and Mikawa, T.
Terminal diversification of the myocyte lineage generates Purkinje fibers of the cardiac conduction system 1423-1431

Grindley, J. C., Davidson, D. R. and Hill, R. E.
The role of *Pax-6* in eye and nasal development 1433-1442

Stier, H. and Schlosshauer, B.
Axonal guidance in the chicken retina 1443-1454

Fletcher, J. C., Burtis, K. C., Hogness, D. S. and Thummel, C. S.
The *Drosophila E74* gene is required for metamorphosis and plays a role in the polytene chromosome puffing response to ecdysone 1455-1465

Danos, M. C. and Yost, H. J.
Linkage of cardiac left-right asymmetry and dorsal-anterior development in *Xenopus* 1467-1474

LaBonne, C., Burke, B. and Whitman, M.
Role of MAP kinase in mesoderm induction and axial patterning during *Xenopus* development 1475-1486

MacGregor, G. R., Zambrowicz, B. P. and Soriano, P.
Tissue non-specific alkaline phosphatase is expressed in both embryonic and extraembryonic lineages during mouse embryogenesis but is not required for migration of primordial germ cells 1487-1496

Wikramanayake, A. H., Brandhorst, B. P. and Klein, W. H.
Autonomous and non-autonomous differentiation of ectoderm in different sea urchin species 1497-1505

Davies, J., Lyon, M., Gallagher, J. and Garrod, D.
Sulphated proteoglycan is required for collecting duct growth and branching but not nephron formation during kidney development 1507-1517

Sessions, R. A. and Zambryski, P. C.
Arabidopsis gynoecium structure in the wild type and *ettin* mutants 1519-1532

Conlon, R. A., Reaume, A. G. and Rossant, J.
Notch1 is required for the coordinate segmentation of somites 1533-1545

Sviderskaya, E. V., Wakeling, W. F. and Bennett, D. C.
A cloned, immortal line of murine melanoblasts inducible to differentiate to melanocytes 1547-1557

Hutter, H. and Schnabel, R.
Specification of anterior-posterior differences within the AB lineage in the *C. elegans* embryo: a polarising induction 1559-1568

ESSAY IN DEVELOPMENT

Slack, J. M. W.
Developmental biology of the pancreas 1569-1580

CONTRIBUTED PAPERS

Kinoshita, K. and Asashima, M.
Effect of activin and lithium on isolated *Xenopus* animal blastomeres and response alteration at the midblastula transition 1581-1589

Yan, L., Pollock, G. H., Nagase, H. and Sarras Jr, M. P.
A 25.7×10^3 M_r hydra metalloproteinase (HMP1), a member of the astacin family, localizes to the extracellular matrix of *Hydra vulgaris* in a head-specific manner and has a developmental function 1591-1602

Hacker, A., Capel, B., Goodfellow, P. and Lovell-Badge, R.
Expression of *Sry*, the mouse sex determining gene 1603-1614

Norrander, J. M., Linck, R. W. and Stephens, R. E.
Transcriptional control of tektin A mRNA correlates with cilia development and length determination during sea urchin embryogenesis 1615-1623

Schwartz, C., Locke, J., Nishida, C. and Kornberg, T. B.
Analysis of *cubitus interruptus* regulation in *Drosophila* embryos and imaginal disks 1625-1635

Sokol, S. Y., Klingensmith, J., Perrimon, N. and Itoh, K.
Dorsalizing and neuralizing properties of Xdsh, a maternally expressed *Xenopus* homolog of *dishevelled* 1637-1647

Tiong, S. Y. K., Nash, D. and Bender, W.
Dorsal wing, a locus that affects dorsoventral wing patterning in *Drosophila* 1649-1656

Masteller, E. L., Larsen, R. D., Carlson, L. M., Pickel, J. M., Nickoloff, B., Lowe, J., Thompson, C. B. and Lee, K. P.
Chicken B cells undergo discrete developmental changes in surface carbohydrate structure that appear to play a role in directing lymphocyte migration during embryogenesis 1657-1667

Reneker, L. W., Silversides, D. W., Patel, K. and Overbeek, P. A.
TGFα can act as a chemoattractant to perioptic mesenchymal cells in developing mouse eyes 1669-1680

Chiang, A., O'Connor, M. B., Paro, R., Simon, J. and Bender, W.
Discrete Polycomb-binding sites in each parasegmental domain of the bithorax complex 1681-1689

Serrano, N., Brock, H. W., Demeret, C., Dura, J.-M., Randsholt, N. B., Kornberg, T. B. and Maschat, F.
polyhomeotic appears to be a target of Engrailed regulation in *Drosophila* 1691-1703

Fontaine-Perus, J., Jarno, V., Fournier Le Ray, C., Li, Z. and Paulin, D.
Mouse chick chimera: a new model to study the in ovo developmental potentialities of mammalian somites 1705-1718

Feng, L. and Heintz, N.
Differentiating neurons activate transcription of the brain lipid-binding protein gene in radial glia through a novel regulatory element 1719-1730

Gardiner, D. M., Blumberg, B., Komine, Y. and Bryant, S. V.
Regulation of *HoxA* expression in developing and regenerating axolotl limbs 1731-1741

Ono, K., Bansal, R., Payne, J., Rutishauser, U. and Miller, R. H.
Early development and dispersal of oligodendrocyte precursors in the embryonic chick spinal cord 1743-1754

Barth, K. A. and Wilson, S. W.
Expression of zebrafish *nk2.2* is influenced by *sonic hedgehog/vertebrate hedgehog-1* and demarcates a zone of neuronal differentiation in the embryonic forebrain 1755-1768

Lundgren, S. E., Callahan, C. A., Thor, S. and Thomas, J. B.
Control of neuronal pathway selection by the *Drosophila* LIM homeodomain gene *apterous* 1769-1773

Robbie, E. P., Peterson, M., Amaya, E. and Musci, T. J.
Temporal regulation of the *Xenopus* FGF receptor in development: a translation inhibitory element in the 3′ untranslated region 1775-1785

Kelly, G. M., Greenstein, P., Erezyilmaz, D. F. and Moon, R. T.
Zebrafish *wnt8* and *wnt8b* share a common activity but are involved in distinct developmental pathways 1787-1799

John, A., Smith, S. T. and Jaynes, J. B.
Inserting the Ftz homeodomain into Engrailed creates a dominant transcriptional repressor that specifically turns off Ftz target genes in vivo 1801-1813

Boyle, M. and DiNardo, S.
Specification, migration and assembly of the somatic cells of the *Drosophila* gonad 1815-1825

Yazaki, I., Tosti, E. and Dale, B.
Cytoskeletal elements link calcium channel activity and the cell cycle in early sea urchin embryos 1827-1831

Stoker, A. W., Gehrig, B., Haj, F. and Bay, B.-H.
Axonal localisation of the CAM-like tyrosine phosphatase CRYPα: a signalling molecule of embryonic growth cones 1833-1844

Dickson, M. C., Martin, J. S., Cousins, F. M., Kulkarni, A. B., Karlsson, S. and Akhurst, R. J.
Defective haematopoiesis and vasculogenesis in transforming growth factor-β1 knock out mice 1845-1854

Casares, F. and Sánchez-Herrero, E.
Regulation of the *infraabdominal* regions of the bithorax complex of *Drosophila* by gap genes 1855-1866

Poirie, M., Niederer, E. and Steinmann-Zwicky, M.
A sex-specific number of germ cells in embryonic gonads of *Drosophila* 1867-1873

Schweisguth, F.
Suppressor of Hairless is required for signal reception during lateral inhibition in the *Drosophila* pupal notum 1875-1884

Pankratz, M. J. and Hoch, M.
Control of epithelial morphogenesis by cell signaling and integrin molecules in the *Drosophila* foregut 1885-1898

Curtis, D., Apfeld, J. and Lehmann, R.
nanos is an evolutionarily conserved organizer of anterior-posterior polarity 1899-1910

Gutknecht, D. R., Koster, C. H., Tertoolen, L. G. J., de Laat, S. W. and Durston, A. J.
Intracellular acidification of gastrula ectoderm is important for posterior axial development in *Xenopus* 1911-1925

Knecht, A. K., Good, P. J., Dawid, I. B. and Harland, R. M.
Dorsal-ventral patterning and differentiation of noggin-induced neural tissue in the absence of mesoderm 1927-1936

Wheatley, S., Kulkarni, S. and Karess, R.
Drosophila nonmuscle myosin II is required for rapid cytoplasmic transport during oogenesis and for axial nuclear migration in early embryos 1937-1946

Epifano, O., Liang, L.-f., Familari, M., Moos Jr., M. C. and Dean, J.
Coordinate expression of the three zona pellucida genes during mouse oogenesis 1947-1956

Makabe, K. W., Kirchhamer, C. V., Britten, R. J. and Davidson, E. H.
Cis-regulatory control of the *SM50* gene, an early marker of skeletogenic lineage specification in the sea urchin embryo 1957-1970

Chan, D. C., Laufer, E., Tabin, C. and Leder, P.
Polydactylous limbs in *Strong's Luxoid* mice result from ectopic polarizing activity 1971-1978

Rushton, E., Drysdale, R., Abmayr, S. M., Michelson, A. M. and Bate, M.
Mutations in a novel gene, *myoblast city*, provide evidence in support of the founder cell hypothesis for *Drosophila* muscle development 1979-1988

Manley, N. R. and Capecchi, M. R.
The role of *Hoxa-3* in mouse thymus and thyroid development 1989-2003

Davidson, L. A., Koehl, M. A. R., Keller, R. and Oster, G. F.
How do sea urchins invaginate? Using biomechanics to distinguish between mechanisms of primary invagination 2005-2018

Jarman, A. P., Sun, Y., Jan, L. Y. and Jan, Y. N.
Role of the proneural gene, *atonal*, in formation of *Drosophila* chordotonal organs and photoreceptors 2019-2030

Engelhardt, J. F., Schlossberg, H., Yankaskas J. R. and Dudus, L.
Progenitor cells of the adult human airway involved in submucosal gland development 2031-2046

Wickramasinghe, D., Becker, S., Ernst, M. K., Resnick, J. L., Centanni, J. M., Tessarollo, L., Grabel, L. B. and Donovan, P. J.
Two CDC25 homologues are differentially expressed during mouse development 2047-2056

Clark, S. E., Running, M. P. and Meyerowitz, E. M.
CLAVATA3 is a specific regulator of shoot and floral meristem development affecting the same processes as *CLAVATA1* 2057-2067

Newman-Smith, E. D. and Werb, Z.
Stem cell defects in parthenogenetic peri-implantation embryos 2069-2077

Robinson, G. W., McKnight, R. A., Smith, G. H. and Hennighausen, L.
Mammary epithelial cells undergo secretory differentiation in cycling virgins but require pregnancy for the establishment of terminal differentiation 2079-2090

Ramírez, F.-A., Wedeen, C. J., Stuart, D. K., Lans, D. and Weisblat, D. A.
Identification of a neurogenic sublineage required for CNS segmentation in an Annelid 2091-2097

Dickinson, M. E., Selleck, M. A. J., McMahon, A. P. and Bronner-Fraser, M.
Dorsalization of the neural tube by the non-neural ectoderm 2099-2106

Maggert, K., Levine, M. and Frasch, M.
The somatic-visceral subdivision of the embryonic mesoderm is initiated by dorsal gradient thresholds in *Drosophila* 2107-2116

González-Crespo, S. and Morata, G.
Control of *Drosophila* adult pattern by *extradenticle* 2117-2125

Kapoun, A. M. and Kaufman, T. C.
A functional analysis of 5′ intronic and promoter regions of the homeotic gene *proboscipedia* in *Drosophila melanogaster* 2127-2141

Waites, R. and Hudson, A.
phantastica: a gene required for dorsoventrality of leaves in *Antirrhinum majus* 2143-2154

Kessler, D. S. and Melton, D. A.
Induction of dorsal mesoderm by soluble, mature Vg1 protein 2155-2164

O'Rourke, N. A., Sullivan, D. P., Kaznowski, C. E., Jacobs, A. A. and McConnell, S. K.
Tangential migration of neurons in the developing cerebral cortex 2165-2176

Cui, Y., Brown, J. D., Moon, R. T. and Christian, J. L.
Xwnt-8b: a maternally expressed *Xenopus Wnt* gene with a potential role in establishing the dorsoventral axis 2177-2186

Gard, A. L., Burrell, M. R., Pfeiffer, S. E., Rudge, J. S. and Williams II, W. C.
Astroglial control of oligodendrocyte survival mediated by PDGF and leukemia inhibitory factor-like protein 2187-2197

Donjacour, A. A. and Cunha, G. R.
Induction of prostatic morphology and secretion in urothelium by seminal vesicle mesenchyme 2199-2207

Galindo, R. L., Edwards, D. N., Gillespie, S. K. H. and Wasserman, S. A.
Interaction of the pelle kinase with the membrane-associated protein tube is required for transduction of the dorsoventral signal in *Drosophila* embryos 2209-2218

Schnabel, R.
Duels without obvious sense: counteracting inductions involved in body wall muscle development in the *Caenorhabditis elegans* embryo 2219-2232

Ayabe, T., Kopf, G. S. and Schultz, R. M.
Regulation of mouse egg activation: presence of ryanodine receptors and effects of microinjected ryanodine and cyclic ADP ribose on uninseminated and inseminated eggs 2233-2244

Grieshammer, U., McGrew, M. J. and Rosenthal, N.
Role of methylation in maintenance of positionally restricted transgene expression in developing muscle 2245-2253

Murphy, A. M., Lee, T., Andrews, C. M., Shilo, B.-Z. and Montell, D. J.
The Breathless FGF receptor homolog, a downstream target of *Drosophila* C/EBP in the developmental control of cell migration 2255-2263

Zecca, M., Basler, K. and Struhl, G.
Sequential organizing activities of engrailed, hedgehog and decapentaplegic in the *Drosophila* wing 2265-2278

Ma, C. and Moses, K.
wingless and *patched* are negative regulators of the morphogenetic furrow and can affect tissue polarity in the developing *Drosophila* compound eye 2279-2289

Cubitt, A. B., Firtel, R. A., Fischer, G., Jaffe, L. F. and Miller, A. L.
Patterns of free calcium in multicellular stages of *Dictyostelium* expressing jellyfish apoaequorin 2291-2301

Landolt, R. M., Vaughan, L., Winterhalter, K. H. and Zimmermann, D. R.
Versican is selectively expressed in embryonic tissues that act as barriers to neural crest cell migration and axon outgrowth 2303-2312

González-Gaitán, M. and Jäckle, H.
Invagination centers within the *Drosophila* stomatogastric nervous system anlage are positioned by *Notch*-mediated signaling which is spatially controlled through *wingless* 2313-2325

Berninger, B., Marty, S., Zafra, F., da Penha Berzaghi, M., Thoenen, H. and Lindholm, D.
GABAergic stimulation switches from enhancing to repressing BDNF expression in rat hippocampal neurons during maturation in vitro 2327-2335

Ekker, S. C., McGrew, L. L., Lai, C.-J., Lee, J. J., von Kessler, D. P., Moon, R. T. and Beachy, P. A.
Distinct expression and shared activities of members of the *hedgehog* gene family of *Xenopus laevis* 2337-2347

Lai, C.-J., Ekker, S. C., Beachy, P. A. and Moon, R. T.
Patterning of the neural ectoderm of *Xenopus laevis* by the amino-terminal product of hedgehog autoproteolytic cleavage 2349-2360

Hall, A. K. and MacPhedran, S. E.
Multiple mechanisms regulate sympathetic neuronal phenotype 2361-2371

Schmidt, E. E. and Schibler, U.
High accumulation of components of the RNA polymerase II transcription machinery in rodent spermatids 2373-2383

Herrup, K. and Busser, J. C.
The induction of multiple cell cycle events precedes target-related neuronal death 2385-2395

Kimura, Y. and Yanagimachi, R.
Mouse oocytes injected with testicular spermatozoa or round spermatids can develop into normal offspring 2397-2405

Bettenhausen, B., Hrabě de Angelis, M., Simon, D., Guénet, J.-L. and Gossler, A.
Transient and restricted expression during mouse embryogenesis of *Dll1*, a murine gene closely related to *Drosophila Delta* 2407-2418

Weil, M., Itin, A. and Keshet, E.
A role for mesenchyme-derived tachykinins in tooth and mammary gland morphogenesis 2419-2428

Cornell, R. A., Musci, T. J. and Kimelman, D.
FGF is a prospective competence factor for early activin-type signals in *Xenopus* mesoderm induction 2429-2437

Gannon, M. and Bader, D.
Initiation of cardiac differentiation occurs in the absence of anterior endoderm 2439-2450

Wehrli, M. and Tomlinson, A.
Epithelial planar polarity in the developing *Drosophila* eye 2451-2459

Biffo, S., Offenhäuser, N., Carter, B. D. and Barde, Y.-A.
Selective binding and internalisation by truncated receptors restrict the availability of BDNF during development 2461-2470

Liu, Z., Kirch, S. and Ambros, V.
The *Caenorhabditis elegans* heterochronic gene pathway controls stage-specific transcription of collagen genes 2471-2478

Harrison, S. M., Dunwoodie, S. L., Arkell, R. M., Lehrach, H. and Beddington, R. S. P.
Isolation of novel tissue-specific genes from cDNA libraries representing the individual tissue constituents of the gastrulating mouse embryo 2479-2489

Rougvie, A. E. and Ambros, V.
The heterochronic gene *lin-29* encodes a zinc finger protein that controls a terminal differentiation event in *Caenorhabditis elegans* 2491-2500

Miller, J., Fraser, S. E. and McClay, D.
Dynamics of thin filopodia during sea urchin gastrulation 2501-2511

Cross, J. C., Flannery, M. L., Blanar, M. A., Steingrimsson, E., Jenkins, N. A., Copeland, N. G., Rutter, W. J. and Werb, Z.
Hxt encodes a basic helix-loop-helix transcription factor that regulates trophoblast cell development 2513-2523

Church, D. L., Guan, K.-L. and Lambie, E. J.
Three genes of the MAP kinase cascade, *mek-2*, *mpk-1/sur-1* and *let-60 ras*, are required for meiotic cell cycle progression in *Caenorhabditis elegans* 2525-2535

Martí, E., Takada, R., Bumcrot, D. A., Sasaki, H. and McMahon, A. P.
Distribution of Sonic hedgehog peptides in the developing chick and mouse embryo 2537-2547

Currie, D. A. and Bate, M.
Innervation is essential for the development and differentiation of a sex-specific adult muscle in *Drosophila melanogaster* 2549-2557

Souza, P., Kuliszewski, M., Wang, J., Tseu, I., Tanswell, A. K. and Post, M.
PDGF-AA and its receptor influence early lung branching via an epithelial–mesenchymal interaction 2559-2567

Trainor, P. A. and Tam, P. P. L.
Cranial paraxial mesoderm and neural crest cells of the mouse embryo: co-distribution in the craniofacial mesenchyme but distinct segregation in branchial arches 2569-2581

Brill, G., Kahane, N., Carmeli, C., von Schack, D., Barde, Y.-A. and Kalcheim, C.
Epithelial-mesenchymal conversion of dermatome progenitors requires neural tube-derived signals: characterization of the role of Neurotrophin-3 2583-2594

Woo, K. and Fraser, S. E.
Order and coherence in the fate map of the zebrafish nervous system 2595-2609

Paululat, A., Burchard, S. and Renkawitz-Pohl, R.
Fusion from myoblasts to myotubes is dependent on the *rolling stone* gene (*rost*) of *Drosophila* 2611-2620

Chien, C.-B., Cornel, E. M. and Holt, C. E.
Absence of topography in precociously innervated tecta 2621-2631

Matsuno, K., Diederich, R. J., Go, M. J., Blaumueller, C. M. and Artavanis-Tsakonas, S.
Deltex acts as a positive regulator of Notch signaling through interactions with the Notch ankyrin repeats 2633-2644

Yue, C., White, K. L., Reed, W. A. and Bunch, T. D.
The existence of inositol 1,4,5-trisphosphate and ryanodine receptors in mature bovine oocytes 2645-2654

Koga, M. and Ohshima, Y.
Mosaic analysis of the *let-23* gene function in vulval induction of *Caenorhabditis elegans* 2655-2666

Andres, A. J. and Thummel, C. S.
The *Drosophila* 63F early puff contains *E63-1*, an ecdysone-inducible gene that encodes a novel Ca^{2+}-binding protein 2667-2679

Heller, S., Finn, T. P., Huber, J., Nishi, R., Geißen, M., Püschel, A. W. and Rohrer, H.
Analysis of function and expression of the chick GPA receptor (GPARα) suggests multiple roles in neuronal development 2681-2693

Fuhrmann, S., Kirsch, M. and Hofmann, H.-D.
Ciliary neurotrophic factor promotes chick photoreceptor development in vitro 2695-2706

Grapin-Botton, A., Bonnin, M.-A., McNaughton, L. A., Krumlauf, R. and Le Douarin, N. M.
Plasticity of transposed rhombomeres: Hox gene induction is correlated with phenotypic modifications 2707-2721

Talbert, P. B., Adler, H. T., Parks, D. W. and Comai, L.
The *REVOLUTA* gene is necessary for apical meristem development and for limiting cell divisions in the leaves and stems of *Arabidopsis thaliana* 2723-2735

Rongo, C., Gavis, E. R. and Lehmann, R.
Localization of *oskar* RNA regulates *oskar* translation and requires Oskar protein 2737-2746

Harada, Y., Yasuo, H. and Satoh, N.
A sea urchin homologue of the chordate *Brachyury* (*T*) gene is expressed in the secondary mesenchyme founder cells 2747-2754

Shih, J. and Fraser, S. E.
Distribution of tissue progenitors within the shield region of the zebrafish gastrula 2755-2765

Villano, J. L. and Katz, F. N.
four-jointed is required for intermediate growth in the proximal-distal axis in *Drosophila* 2767-2777

Moser, M., Imhof, A., Pscherer, A., Bauer, R., Amselgruber, W., Sinowatz, F., Hofstädter, F., Schüle, R. and Buettner, R.
Cloning and characterization of a second AP-2 transcription factor: AP-2β 2779-2788

Holowacz, T. and Elinson, R. P.
Properties of the dorsal activity found in the vegetal cortical cytoplasm of *Xenopus* eggs 2789-2798

McCormick, A., Coré, N., Kerridge, S. and Scott, M. P.
Homeotic response elements are tightly linked to tissue-specific elements in a transcriptional enhancer of the *teashirt* gene 2799-2812

Rulifson, E. J. and Blair, S. S.
Notch regulates *wingless* expression and is not required for reception of the paracrine *wingless* signal during wing margin neurogenesis in *Drosophila* 2813-2824

Kerk, N. M. and Feldman, L. J.
A biochemical model for the initiation and maintenance of the quiescent center: implications for organization of root meristems 2825-2833

Treisman, J. E., Lai, Z.-C. and Rubin, G. M.
shortsighted acts in the *decapentaplegic* pathway in *Drosophila* eye development and has homology to a mouse TGF-β-responsive gene 2835-2845

Müller, J., Gaunt, S. and Lawrence, P. A.
Function of the Polycomb protein is conserved in mice and flies 2847-2852

Weng, A., Magnuson, T. and Storb, U.
Strain-specific transgene methylation occurs early in mouse development and can be recapitulated in embryonic stem cells 2853-2859

Zachgo, S., Silva, E. de A., Motte, P., Tröbner, W., Saedler, H. and Schwarz-Sommer, Z.
Functional analysis of the *Antirrhinum* floral homeotic *DEFICIENS* gene in vivo and in vitro by using a temperature-sensitive mutant 2861-2875

Miller III, D. M. and Niemeyer, C. J.
Expression of the *unc-4* homeoprotein in *Caenorhabditis elegans* motor neurons specifies presynaptic input 2877-2886

Day, C. D., Galgoci, B. F. C. and Irish, V. F.
Genetic ablation of petal and stamen primordia to elucidate cell interactions during floral development 2887-2895

Niranjan, B., Buluwela, L., Yant, J., Perusinghe, N., Atherton, A., Phippard, D., Dale, T., Gusterson, B. and Kamalati, T.
HGF/SF: a potent cytokine for mammary growth, morphogenesis and development 2897-2908

Dear, T. N., Colledge, W. H., Carlton, M. B. L., Lavenir, I., Larson, T., Smith, A. J. H., Warren, A. J., Evans, M. J., Sofroniew, M. V. and Rabbitts, T. H.
The *Hox11* gene is essential for cell survival during spleen development 2909-2915

Yamada, G., Mansouri, A., Torres, M., Stuart, E. T., Blum, M., Schultz, M., De Robertis, E. M. and Gruss, P.
Targeted mutation of the murine *goosecoid* gene results in craniofacial defects and neonatal death 2917-2922

Brewster, R. and Bodmer, R.
Origin and specification of type II sensory neurons in *Drosophila* 2923-2936

McKearin, D. and Ohlstein, B.
A role for the *Drosophila* Bag-of-marbles protein in the differentiation of cystoblasts from germline stem cells 2937-2947

Worrad, D. M., Turner, B. M. and Schultz, R. M.
Temporally restricted spatial localization of acetylated isoforms of histone H4 and RNA polymerase II in the 2-cell mouse embryo 2949-2959

Strome, S., Martin, P., Schierenberg, E. and Paulsen, J.
Transformation of the germ line into muscle in *mes-1* mutant embryos of *Caenorhabditis elegans* 2961-2972

Castelli-Gair, J. and Akam, M.
How the Hox gene *Ultrabithorax* specifies two different segments: the significance of spatial and temporal regulation within metameres 2973-2982

Griffin, K., Patient, R. and Holder, N.
Analysis of FGF function in normal and *no tail* zebrafish embryos reveals separate mechanisms for formation of the trunk and the tail 2983-2994

Kuwabara, P. E. and Kimble, J.
A predicted membrane protein, TRA-2A, directs hermaphrodite development in *Caenorhabditis elegans* 2995-3004

Rivera-Pérez, J. A., Mallo, M., Gendron-Maguire, M., Gridley, T. and Behringer, R. R.
goosecoid is not an essential component of the mouse gastrula organizer but is required for craniofacial and rib development 3005-3012

Serano, T. L. and Cohen, R. S.
Gratuitous mRNA localization in the *Drosophila* oocyte 3013-3021

Rittenhouse, K. R. and Berg, C. A.
Mutations in the *Drosophila* gene *bullwinkle* cause the formation of abnormal eggshell structures and bicaudal embryos 3023-3033

Bissen, S. T.
Expression of the cell cycle control gene, *cdc25*, is constitutive in the segmental founder cells but is cell-cycle-regulated in the micromeres of leech embryos 3035-3043

Zheng, L., Zhang, J. and Carthew, R. W.
frizzled regulates mirror-symmetric pattern formation in the *Drosophila* eye 3045-3055

Serra, R. and Moses, H. L.
pRb is necessary for inhibition of N-*myc* expression by TGF-β1 in embryonic lung organ cultures 3057-3066

Margolis, J. S., Borowsky, M. L., Steingrímsson, E., Shim, C. W., Lengyel, J. A. and Posakony, J. W.
Posterior stripe expression of *hunchback* is driven from two promoters by a common enhancer element 3067-3077

Taylor, B. J. and Knittel, L. M.
Sex-specific differentiation of a male-specific abdominal muscle, the Muscle of Lawrence, is abnormal in hydroxyurea-treated and in *fruitless* male flies 3079-3088

Charlton, W. L., Keen, C. L., Merriman, C., Lynch, P., Greenland, A. J. and Dickinson, H. G.
Endosperm development in *Zea mays*; implication of gametic imprinting and paternal excess in regulation of transfer layer development 3089-3097

Ataliotis, P., Symes, K., Chou, M. M., Ho, L. and Mercola, M.
PDGF signalling is required for gastrulation of *Xenopus laevis* 3099-3110

Vervoort, M., Zink, D., Pujol, N., Victoir, K., Dumont, N., Ghysen, A. and Dambly-Chaudière, C.
Genetic determinants of sense organ identity in *Drosophila*: regulatory interactions between *cut* and *poxn* 3111-3120

Kengaku, M. and Okamoto, H.
bFGF as a possible morphogen for the anteroposterior axis of the central nervous system in *Xenopus* 3121-3130

Moore, S. W., Keller, R. E. and Koehl, M. A. R.
The dorsal involuting marginal zone stiffens anisotropically during its convergent extension in the gastrula of *Xenopus laevis* 3131-3140

Stainier, D. Y. R., Weinstein, B. M., Detrich III, H. W., Zon, L. I. and Fishman, M. C.
cloche, an early acting zebrafish gene, is required by both the endothelial and hematopoietic lineages 3141-3150

Chan, D. C., Wynshaw-Boris, A. and Leder, P.
Formin isoforms are differentially expressed in the mouse embryo and are required for normal expression of *fgf-4* and *shh* in the limb bud 3151-3162

Roberts, D. J., Johnson, R. L., Burke, A. C., Nelson, C. E., Morgan, B. A. and Tabin, C.
Sonic hedgehog is an endodermal signal inducing *Bmp-4* and *Hox* genes during induction and regionalization of the chick hindgut 3163-3174

Martindale, M. Q. and Henry, J. Q.
Modifications of cell fate specification in equal-cleaving nemertean embryos: alternate patterns of spiralian development 3175-3185

Spana, E. P. and Doe, C. Q.
The prospero transcription factor is asymmetrically localized to the cell cortex during neuroblast mitosis in *Drosophila* 3187-3195

Katsuyama, Y., Wada, S., Yasugi, S. and Saiga, H.
Expression of the *labial* group Hox gene *HrHox-1* and its alteration induced by retinoic acid in development of the ascidian *Halocynthia roretzi* 3197-3205

Qiao, J., Cohen, D. and Herzlinger, D.
The metanephric blastema differentiates into collecting system and nephron epithelia in vitro 3207-3214

Ransick, A. and Davidson, E. H.
Micromeres are required for normal vegetal plate specification in sea urchin embryos 3215-3222

Millar, S. E., Miller, M. W., Stevens, M. E. and Barsh, G. S.
Expression and transgenic studies of the mouse *agouti* gene provide insight into the mechanisms by which mammalian coat color patterns are generated 3223-3232

Cui, X. and Doe, C. Q.
The role of the cell cycle and cytokinesis in regulating neuroblast sublineage gene expression in the *Drosophila* CNS 3233-3243

Bashaw, G. J. and Baker, B. S.
The *msl-2* dosage compensation gene of *Drosophila* encodes a putative DNA-binding protein whose expression is sex specifically regulated by *Sex-lethal* 3245-3258

Jones, K. T., Carroll, J., Merriman, J. A., Whittingham, D. G. and Kono, T.
Repetitive sperm-induced Ca^{2+} transients in mouse oocytes are cell cycle dependent 3259-3266

Macdonald, R., Barth, K. A., Xu, Q., Holder, N., Mikkola, I. and Wilson, S. W.
Midline signalling is required for Pax gene regulation and patterning of the eyes 3267-3278

Acampora, D., Mazan, S., Lallemand, Y., Avantaggiato, V., Maury, M., Simeone, A. and Brûlet, P.
Forebrain and midbrain regions are deleted in *Otx2*$^{-/-}$ mutants due to a defective anterior neuroectoderm specification during gastrulation 3279-3290

Oka, C., Nakano, T., Wakeham, A., Luis de la Pompa, J., Mori, C., Sakai, T., Okazaki, S., Kawaichi, M., Shiota, K., Mak, T. W. and Honjo, T.
Disruption of the mouse *RBP-Jκ* gene results in early embryonic death 3291-3301

Laskowski, M. J., Williams, M. E., Nusbaum, H. C. and Sussex, I. M.
Formation of lateral root meristems is a two-stage process 3303-3310

Domingo, C. and Keller, R.
Induction of notochord cell intercalation behavior and differentiation by progressive signals in the gastrula of *Xenopus laevis* 3311-3321

Hsu, D. R., Chuang, P.-T. and Meyer, B. J.
DPY-30, a nuclear protein essential early in embryogenesis for *Caenorhabditis elegans* dosage compensation 3323-3334

Guimarães, M. J., Bazan, J. F., Zlotnik, A., Wiles, M. V., Grimaldi, J. C., Lee, F. and McClanahan, T.
A new approach to the study of haematopoietic development in the yolk sac and embryoid bodies 3335-3346

Patapoutian, A., Yoon, J. K., Miner, J. H., Wang, S., Stark, K. and Wold, B.
Disruption of the mouse MRF4 gene identifies multiple waves of myogenesis in the myotome 3347-3358

Tabata, T., Schwartz, C., Gustavson, E., Ali, Z. and Kornberg, T. B.
Creating a *Drosophila* wing de novo, the role of *engrailed*, and the compartment border hypothesis 3359-3369

Richardson, H., O'Keefe, L. V., Marty, T. and Saint, R.
Ectopic cyclin E expression induces premature entry into S phase and disrupts pattern formation in the *Drosophila* eye imaginal disc 3371-3379

Schimmang, T., Minichiello, L., Vazquez, E., San Jose, I., Giraldez, F., Klein, R. and Represa, J.
Developing inner ear sensory neurons require TrkB and TrkC receptors for innervation of their peripheral targets 3381-3391

Staehling-Hampton, K., Laughon, A. S. and Hoffmann, F. M.
A *Drosophila* protein related to the human zinc finger transcription factor PRDII/MBPI/HIV-EP1 is required for *dpp* signaling 3393-3403

de Celis, J. F., Llimargas, M. and Casanova, J.
ventral veinless, the gene encoding the Cf1a transcription factor, links positional information and cell differentiation during embryonic and imaginal development in *Drosophila melanogaster* 3405-3416

Hutter, H. and Schnabel, R.
Establishment of left-right asymmetry in the *Caenorhabditis elegans* embryo: a multistep process involving a series of inductive events 3417-3424

Thompson, E. M., Legouy, E., Christians, E. and Renard, J.-P.
Progressive maturation of chromatin structure regulates *HSP70.1* gene expression in the preimplantation mouse embryo 3425-3437

Yin, X., Watanabe, M. and Rutishauser, U.
Effect of polysialic acid on the behavior of retinal ganglion cell axons during growth into the optic tract and tectum 3439-3446

Guillén, I., Mullor, J. L., Capdevila, J., Sánchez-Herrero, E., Morata, G. and Guerrero, I.
The function of *engrailed* and the specification of *Drosophila* wing pattern 3447-3456

Roegiers, F., McDougall, A. and Sardet, C.
The sperm entry point defines the orientation of the calcium-induced contraction wave that directs the first phase of cytoplasmic reorganization in the ascidian egg 3457-3466

de Celis, J. F. and Ruiz-Gómez, M.
groucho and *hedgehog* regulate *engrailed* expression in the anterior compartment of the *Drosophila* wing 3467-3476

Eberhart, C. G. and Wasserman, S. A.
The *pelota* locus encodes a protein required for meiotic cell division: an analysis of G_2/M arrest in *Drosophila* spermatogenesis 3477-3486

Erratum to sequence published in *Development* **121**,1637-1647. 3487

Spana, E. P., Kopczynski, C., Goodman, C. S. and Doe, C. Q.
Asymmetric localization of numb autonomously determines sibling neuron identity in the *Drosophila* CNS 3489-3494

Jaglarz, M. K. and Howard, K. R.
The active migration of *Drosophila* primordial germ cells 3495-3503

Vodicka, M. A. and Gerhart, J. C.
Blastomere derivation and domains of gene expression in the Spemann Organizer of *Xenopus laevis* 3505-3518

Treisman, J. E. and Rubin, G. M.
wingless inhibits morphogenetic furrow movement in the *Drosophila* eye disc 3519-3527

Haegel, H., Larue, L., Ohsugi, M., Fedorov, L., Herrenknecht, K. and Kemler, R.
Lack of β-catenin affects mouse development at gastrulation 3529-3537

Wysolmerski, J. J., McCaughern-Carucci, J. F., Daifotis, A. G., Broadus, A. E. and Philbrick, W. M.
Overexpression of parathyroid hormone-related protein or parathyroid hormone in transgenic mice impairs branching morphogenesis during mammary gland development 3539-3547

Emery, J. F. and Bier, E.
Specificity of CNS and PNS regulatory subelements comprising pan-neural enhancers of the *deadpan* and *scratch* genes is achieved by repression 3549-3560

Royet, J. and Finkelstein, R.
Pattern formation in *Drosophila* head development: the role of the *orthodenticle* homeobox gene 3561-3572

Roseman, R. R., Swan, J. M. and Geyer, P. K.
A *Drosophila* insulator protein facilitates dosage compensation of the X chromosome *mini-white* gene located at autosomal insertion sites 3573-3582

Rosenbaum, T., Boissy, Y. L., Kombrinck, K., Brannan, C. I., Jenkins, N. A., Copeland, N. G. and Ratner, N.
Neurofibromin-deficient fibroblasts fail to form perineurium in vitro 3583-3592

Bradshaw, A. D., McNagny, K. M., Gervin, D. B., Cann, G. M., Graf, T. and Clegg, D. O.
Integrin $\alpha_2\beta_1$ mediates interactions between developing embryonic retinal cells and collagen 3593-3602

MacArthur, C. A., Lawshé, A., Xu, J., Santos-Ocampo, S., Heikinheimo, M., Chellaiah, A. T. and Ornitz, D. M.
FGF-8 isoforms activate receptor splice forms that are expressed in mesenchymal regions of mouse development 3603-3613

Chow, K. L., Hall, D. H. and Emmons, S. W.
The *mab-21* gene of *Caenorhabditis elegans* encodes a novel protein required for choice of alternate cell fates 3615-3626

Lamb, T. M. and Harland, R. M.
Fibroblast growth factor is a direct neural inducer, which combined with noggin generates anterior-posterior neural pattern 3627-3636

Austin, C. P., Feldman, D. E., Ida Jr., J. A. and Cepko, C. L.
Vertebrate retinal ganglion cells are selected from competent progenitors by the action of *Notch* 3637-3650

Jones, C. M., Kuehn, M. R., Hogan, B. L. M., Smith, J. C. and Wright, C. V. E.
Nodal-related signals induce axial mesoderm and dorsalize mesoderm during gastrulation 3651-3662

Rauskolb, C., Smith, K. M., Peifer, M. and Wieschaus, E.
extradenticle determines segmental identities throughout *Drosophila* development 3663-3673

Stern, H. M., Brown, A. M. C. and Hauschka, S. D.
Myogenesis in paraxial mesoderm: preferential induction by dorsal neural tube and by cells expressing *Wnt-1* 3675-3686

Nakato, H., Futch, T. A. and Selleck, S. B.
The *division abnormally delayed (dally)* gene: a putative integral membrane proteoglycan required for cell division patterning during postembryonic development of the nervous system in *Drosophila* 3687-3702

Hidalgo, A., Urban, J. and Brand, A. H.
Targeted ablation of glia disrupts axon tract formation in the *Drosophila* CNS 3703-3712

Weigmann, K. and Lehner, C. F.
Cell fate specification by *even-skipped* expression in the *Drosophila* nervous system is coupled to cell cycle progression 3713-3721

Markussen, F.-H., Michon, A.-M., Breitwieser, W. and Ephrussi, A.
Translational control of *oskar* generates Short OSK, the isoform that induces pole plasm assembly 3723-3732

Krull, C. E., Collazo, A., Fraser, S. E. and Bronner-Fraser, M.
Segmental migration of trunk neural crest: time-lapse analysis reveals a role for PNA-binding molecules 3733-3743

Jennings, B., de Celis, J., Delidakis, C., Preiss, A. and Bray, S.
Role of *Notch* and *achaete-scute* complex in the expression of *Enhancer of split* bHLH proteins 3745-3752

Viviano, C. M., Horton, C. E., Maden, M. and Brockes, J. P.
Synthesis and release of 9-*cis* retinoic acid by the urodele wound epidermis 3753-3762

Yamagata, M. and Sanes, J. R.
Target-independent diversification and target-specific projection of chemically defined retinal ganglion cell subsets 3763-3776

Kelley, M. W., Turner, J. K. and Reh, T. A.
Ligands of steroid/thyroid receptors induce cone photoreceptors in vertebrate retina 3777-3785

Kapur, R. P., Sweetser, D. A., Doggett, B., Siebert, J. R. and Palmiter, R. D.
Intercellular signals downstream of endothelin receptor-B mediate colonization of the large intestine by enteric neuroblasts 3787-3795

Margolis, J. and Spradling, A.
Identification and behavior of epithelial stem cells in the *Drosophila* ovary 3797-3807

Serano, T. L. and Cohen, R. S.
A small predicted stem-loop structure mediates oocyte localization of *Drosophila K10* mRNA 3809-3818

Curtiss, J. and Heilig, J. S.
Establishment of *Drosophila* imaginal precursor cells is controlled by the *Arrowhead* gene 3819-3828

Baylies, M. K., Martinez Arias, A. and Bate, M.
wingless is required for the formation of a subset of muscle founder cells during *Drosophila* embryogenesis 3829-3837

Zaraisky, A. G., Ecochard, V., Kazanskaya, O. V., Lukyanov, S. A., Fesenko, I. V. and Duprat, A.-M.
The homeobox-containing gene *XANF-1* may control development of the Spemann organizer 3839-3847

Therianos, S., Leuzinger, S., Hirth, F., Goodman, C. S. and Reichert, H.
Embryonic development of the *Drosophila* brain: formation of commissural and descending pathways 3849-3860

Heuer, J. G., Li, K. and Kaufman, T. C.
The *Drosophila* homeotic target gene *centrosomin* (*cnn*) encodes a novel centrosomal protein with leucine zippers and maps to a genomic region required for midgut morphogenesis 3861-3876

Soudais, C., Bielinska, M., Heikinheimo, M., MacArthur, C. A., Narita, N., Saffitz, J. E., Simon, M. C., Leiden, J. M. and Wilson, D. B.
Targeted mutagenesis of the transcription factor GATA-4 gene in mouse embryonic stem cells disrupts visceral endoderm differentiation in vitro 3877-3888

Evans, S. M., Yan, W., Murillo, M. P., Ponce, J. and Papalopulu, N.
tinman, a *Drosophila* homeobox gene required for heart and visceral mesoderm specification, may be represented by a family of genes in vertebrates: *XNkx-2.3*, a second vertebrate homologue of *tinman* 3889-3899

Chiang, C., Young, K. E. and Beachy, P. A.
Control of *Drosophila* tracheal branching by the novel homeodomain gene *unplugged*, a regulatory target for genes of the bithorax complex 3901-3912

Hodgetts, R. B., Clark, W. C., O'Keefe, S. L., Schouls, M., Crossgrove, K., Guild, G. M. and von Kalm, L.
Hormonal induction of Dopa decarboxylase in the epidermis of *Drosophila* is mediated by the *Broad-Complex* 3913-3922

ESSAY IN DEVELOPMENT

Shimamura, K., Hartigan, D. J., Martinez, S., Puelles, L. and Rubenstein, J. L. R.
Longitudinal organization of the anterior neural plate and neural tube 3923-3933

CONTRIBUTED PAPERS

Millen, K. J., Hui, C.-C. and Joyner, A. L.
A role for *En-2* and other murine homologues of *Drosophila* segment polarity genes in regulating positional information in the developing cerebellum 3935-3945

Rogge, R., Green, P. J., Urano, J., Horn-Saban, S., Mlodzik, M., Shilo, B.-Z., Hartenstein, V. and Banerjee, U.
The role of *yan* in mediating the choice between cell division and differentiation 3947-3958

Robinson, M. L., MacMillan-Crow, L. A., Thompson, J. A. and Overbeek, P. A.
Expression of a truncated FGF receptor results in defective lens development in transgenic mice 3959-3967

Barnett, S. C., Rosario, M., Doyle, A., Kilbey, A., Lovatt, A. and Gillespie, D. A. F.
Differential regulation of AP-1 and novel TRE-specific DNA-binding complexes during differentiation of oligodendrocyte-type-2-astrocyte (O-2A) progenitor cells 3969-3977

Itoh, K., Tang, T. L., Neel, B. G. and Sokol, S. Y.
Specific modulation of ectodermal cell fates in *Xenopus* embryos by glycogen synthase kinase 3979-3988

Broadus, J. and Doe, C. Q.
Evolution of neuroblast identity: *seven-up* and *prospero* expression reveal homologous and divergent neuroblast fates in *Drosophila* and *Schistocerca* 3989-3996

Sucov, H. M., Izpisúa-Belmonte, J.-C., Gañan, Y. and Evans, R. M.
Mouse embryos lacking RXRα are resistant to retinoic-acid-induced limb defects 3997-4003

Xu, Q., Alldus, G., Holder, N. and Wilkinson, D. G.
Expression of truncated *Sek-1* receptor tyrosine kinase disrupts the segmental restriction of gene expression in the *Xenopus* and zebrafish hindbrain 4005-4016

Hilfiker, A., Amrein, H., Dübendorfer, A., Schneiter, R. and Nöthiger, R.
The gene *virilizer* is required for female-specific splicing controlled by *Sxl*, the master gene for sexual development in *Drosophila* 4017-4026

Grens, A., Mason, E., Marsh, J. L. and Bode, H. R.
Evolutionary conservation of a cell fate specification gene: the *Hydra achaete-scute* homolog has proneural activity in *Drosophila* 4027-4035

Manoukian, A. S., Yoffe, K. B., Wilder, E. L. and Perrimon, N.
The *porcupine* gene is required for *wingless* autoregulation in *Drosophila* 4037-4044

Oliver, G., Mailhos, A., Wehr, R., Copeland, N. G., Jenkins, N. A. and Gruss, P.
Six3, a murine homologue of the *sine oculis* gene, demarcates the most anterior border of the developing neural plate and is expressed during eye development 4045-4055

Torres, M., Gómez-Pardo, E., Dressler, G. R. and Gruss, P.
Pax-2 controls multiple steps of urogenital development 4057-4065

Minichiello, L., Piehl, F., Vazquez, E., Schimmang, T., Hökfelt, T., Represa, J. and Klein, R.
Differential effects of combined *trk* receptor mutations on dorsal root ganglion and inner ear sensory neurons 4067-4075

Tajbakhsh, S. and Buckingham, M. E.
Lineage restriction of the myogenic conversion factor *myf-5* in the brain 4077-4083

Chanut, F. and Heberlein, U.
Role of the morphogenetic furrow in establishing polarity in the *Drosophila* eye 4085-4094

Krasnow, R. E., Wong, L. L. and Adler, P. N.
dishevelled is a component of the *frizzled* signaling pathway in *Drosophila* 4095-4102

Sechrist, J., Nieto, M. A., Zamanian, R. T. and Bronner-Fraser, M.
Regulative response of the cranial neural tube after neural fold ablation: spatiotemporal nature of neural crest regeneration and up-regulation of *Slug* 4103-4115

Appel, B., Korzh, V., Glasgow, E., Thor, S., Edlund, T., Dawid, I. B. and Eisen, J. S.
Motoneuron fate specification revealed by patterned LIM homeobox gene expression in embryonic zebrafish 4117-4125

Smith, C. L., Giordano, H., Schwartz, M. and DeLotto, R.
Spatial regulation of *Drosophila* Snake protease activity in the generation of dorsal-ventral polarity 4127-4135

Behrendtsen, O., Alexander, C. M. and Werb, Z.
Cooperative interactions between extracellular matrix, integrins and parathyroid hormone-related peptide regulate parietal endoderm differentiation in mouse embryos 4137-4148

Weaver, T. A. and White, R. A. H.
headcase, an imaginal specific gene required for adult morphogenesis in *Drosophila melanogaster* 4149-4160

Johnson, R. L., Grenier, J. K. and Scott, M. P.
patched overexpression alters wing disc size and pattern: transcriptional and post-transcriptional effects on *hedgehog* targets 4161-4170

Baima, S., Nobili, F., Sessa, G., Lucchetti, S., Ruberti, I. and Morelli, G.
The expression of the *Athb-8* homeobox gene is restricted to provascular cells in *Arabidopsis thaliana* 4171-4182

Dunin Borkowski, O. M., Brown, N. H. and Bate, M.
Anterior-posterior subdivision and the diversification of the mesoderm in *Drosophila* 4183-4193

Sasaki, H., Ferguson-Smith, A. C., Shum, A. S. W., Barton, S. C. and Surani, M. A.
Temporal and spatial regulation of *H19* imprinting in normal and uniparental mouse embryos 4195-4202

Schultheiss, T. M., Xydas, S. and Lassar, A. B.
Induction of avian cardiac myogenesis by anterior endoderm 4203-4214

Diaz-Benjumea, F. J. and Cohen, S. M.
Serrate signals through Notch to establish a Wingless-dependent organizer at the dorsal/ventral compartment boundary of the *Drosophila* wing 4215-4225

Wolff, C., Sommer, R., Schröder, R., Glaser, G. and Tautz, D.
Conserved and divergent expression aspects of the *Drosophila* segmentation gene *hunchback* in the short germ band embryo of the flour beetle *Tribolium* 4227-4236

Haramis, A. G., Brown, J. M. and Zeller, R.
The *limb deformity* mutation disrupts the SHH/FGF-4 feedback loop and regulation of 5′*HoxD* genes during limb pattern formation 4237-4245

Strutt, D. I. and Mlodzik, M.
Ommatidial polarity in the *Drosophila* eye is determined by the direction of furrow progression and local interactions 4247-4256

Halpern, M. E., Thisse, C., Ho, R. K., Thisse, B., Riggleman, B., Trevarrow, B., Weinberg, E. S., Postlethwait, J. H. and Kimmel, C. B.
Cell-autonomous shift from axial to paraxial mesodermal development in zebrafish *floating head* mutants 4257-4264

Scherer, S. S., Xu, Y.-t., Bannerman, P. G. C., Sherman, D. L. and Brophy, P. J.
Periaxin expression in myelinating Schwann cells: modulation by axon-glial interactions and polarized localization during development 4265-4273

Fitzgerald, K. and Greenwald, I.
Interchangeability of *Caenorhabditis elegans* DSL proteins and intrinsic signalling activity of their extracellular domains in vivo 4275-4282

Holland, P. W. H., Koschorz, B., Holland, L. Z. and Herrmann, B. G.
Conservation of *Brachyury* (*T*) genes in amphioxus and vertebrates: developmental and evolutionary implications 4283-4291

Moos Jr, M., Wang, S. and Krinks, M.
Anti-Dorsalizing Morphogenetic Protein is a novel TGF-β homolog expressed in the Spemann organizer 4293-4301

Lawrence, P. A., Bodmer, R. and Vincent, J.-P.
Segmental patterning of heart precursors in *Drosophila* 4303-4308

Kitsukawa, T., Shimono, A., Kawakami, A., Kondoh, H. and Fujisawa, H.
Overexpression of a membrane protein, neuropilin, in chimeric mice causes anomalies in the cardiovascular system, nervous system and limbs 4309-4318

Schmidt, J., Francois, V., Bier, E. and Kimelman, D.
Drosophila short gastrulation induces an ectopic axis in *Xenopus*: evidence for conserved mechanisms of dorsal-ventral patterning 4319-4328

Hardy, A., Richardson, M. K., Francis-West, P. H., Rodriguez, C., Izpisúa-Belmonte, J.-C., Duprez, D. and Wolpert, L.
Gene expression, polarising activity and skeletal patterning in reaggregated hind limb mesenchyme 4329-4337

Shashikant, C. S., Bieberich, C. J., Belting, H.-G., Wang, J. C. H., Borbély, M. A. and Ruddle, F. H.
Regulation of *Hoxc-8* during mouse embryonic development: identification and characterization of critical elements involved in early neural tube expression 4339-4347

Cox, Wm. G. and Hemmati-Brivanlou, A.
Caudalization of neural fate by tissue recombination and bFGF 4349-4358

Kumar, J. P. and Ready, D. F.
Rhodopsin plays an essential structural role in *Drosophila* photoreceptor development 4359-4370

Fujioka, M., Jaynes, J. B. and Goto, T.
Early *even-skipped* stripes act as morphogenetic gradients at the single cell level to establish *engrailed* expression 4371-4382

Chow, R. L., Roux, G. D., Roghani, M., Palmer, M. A., Rifkin, D. B., Moscatelli, D. A. and Lang, R. A.
FGF suppresses apoptosis and induces differentiation of fibre cells in the mouse lens 4383-4393

Index of Authors and Titles 4395-4407

Subject Index 4409-4447